SOCIAL SCIENCES

Public Health Reports

VOLUME 66 DECEMBER 28, 1951 NUMBER 52

[To be published monthly in 1952—see back cover and page 1738]

IN THIS ISSUE

Rat Ectoparasites and Typhus Fever in Texas

Warfarin as a Plague Suppressive in Hawaii

Warfarin Tests on the Hawaiian Rat

FEDERAL SECURITY AGENCY

PUBLIC HEALTH SERVICE

FEDERAL SECURITY AGENCY
Oscar R. Ewing, Administrator

PUBLIC HEALTH SERVICE
Leonard A. Scheele, Surgeon General

Division of Public Health Methods
G. St. J. Perrott, Chief of Division

CONTENTS

	Page
Observations on rat ectoparasites and typhus fever in San Antonio, Tex. David E. Davis	1717
Use of warfarin-treated oats as a plague suppressive measure in Hawaii. Bertram Gross, Robert H. Baker, and David D. Bonnet	1727
Cage tests with warfarin on the Hawaiian rat, *Rattus hawaiiensis* Stone, and the house mouse, *Mus musculus* Linn., in Hawaii. David D. Bonnet, Edward S. C. Mau, and Bertram Gross	1734
INCIDENCE OF DISEASE	
United States:	
Summary of reports from States	1738
Table of reported cases of communicable diseases	1741
Foreign reports:	
Canada—Provinces—Week ended November 24, 1951	1743
Madagascar—September 1951	1743
Cholera	1744
Plague	1744
Smallpox	1744
Yellow fever	1744

Public Health Reports

Vol. 66 • DECEMBER 28, 1951 • *No. 52*

Observations on Rat Ectoparasites and Typhus Fever in San Antonio, Texas

By David E. Davis*

Studies on the control of typhus fever conducted in San Antonio, Tex., found that rats containing complement-fixing antibodies for typhus fever were widely distributed in the city (*1, 2*). DDT dust to reduce rat fleas was used and resulted in a reduction of human cases of typhus fever (*3, 4*). Observations on rats and typhus fever (*5*) showed that 35 percent of the roof rats and 51 percent of the Norway rats contained typhus complement-fixing antibodies.

Observations on the ectoparasites are presented to make data available on abundance of ectoparasites in the southwestern portions of the area where typhus is endemic. The recent decline in typhus fever (*6*) makes a comparison possible of the data on fleas for the years 1944–45 of high typhus fever rates with data for recent years. These data apparently are the most extensive collected before the reduction in typhus began. Hill and Ingraham (*7*) presented data for the periods of high typhus rates for rural Alabama.

Observations on Fleas

Several species of fleas are found on rats in San Antonio. *Xenopsylla cheopis*, the Indian rat flea, is the most common species and is found wherever there are rats. *Leptopsylla segnis*, the so-called mouse flea, is regularly present and *Echidnophaga gallinacea*, the chicken flea, is common on rats caught in poultry houses and chicken coops. *Ctenocephalides* spp., the cat and dog fleas, are frequently found on rats, especially where cats or dogs are present. *Nosopsyllus fasciatus*, the northern rat flea, is rare. *Pulex irritans*, the human flea, is occasionally present.

Ctenocephalides sp. is the only flea in San Antonio which commonly becomes a pest to humans. This species is the cause of complaints

*Johns Hopkins School of Hygiene and Public Health, Baltimore 5, Md.

that fleas are regularly found in houses and yards. One infestation of *P. irritans* in a shed was observed.

Changes in numbers of fleas found in relation to the seasonal occurrence of typhus fever are of interest. Throughout the southern United States, murine typhus is known to reach a maximum in the summer, shortly after the greatest abundance of rat fleas. For a study of seasonal changes in abundance of fleas in San Antonio, rats were collected systematically throughout the year. The rats were captured in number 0 steel traps, placed alive in a bag, and chloroformed in the laboratory. The rats were carefully combed to obtain all ectoparasites. The ectoparasites were identified under a binocular dissecting scope and then preserved in alcohol.

The rats were collected principally from the southern part of the city. The houses and stores are one-story frame structures. The stores are generally corner groceries. Rats caught in usual places (barns, for example) are not included. An attempt was made to have the sample of rats as comparable as possible from season to season. The collections are probably as random a sample as is practicable. Both Norway rats (*Rattus norvegicus*) and roof rats (*R. rattus*) are represented.

The rats are divided into adult and immature, according to reproductive state (*5*). The indices are calculated according to mean number of fleas per rat and to percent of rats infested as recommended (*8*). Indices are not recorded when less than five rats are in a category.

Table 1 shows that seasonal changes in abundance of *X. cheopis*

Table 1. *Seasonal changes in abundance of* **Xenopsylla cheopis** *(1944–45)*

	May–June			July–August			September–October			November–December		
	Rats	Mean[1]	Percent[2]	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	4	------	------	27	3.5	67	15	3.4	66	6	1.3	66
Immature	8	8.1	88	31	9.0	77	29	1.3	48	26	2.5	42
Norway rats:												
Adults	27	14.1	96	43	11.5	95	22	2.7	73	12	5.1	66
Immature	13	13.9	100	13	10.9	100	17	5.0	88	2	------	------
	December–January–February			March–April			May–June			July–August		
	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Reof rats:												
Adults	53	0.7	34	62	1.3	69	22	3.2	59	19	7.2	68
Immature	58	.9	38	77	1.1	45	63	3.8	70	45	8.4	49
Norway rats:												
Adults	27	1.8	41	60	5.1	58	19	14.7	79	57	8.5	74
Immature	5	6.6	80	34	5.4	65	16	10.0	56	40	8.6	82

[1] Mean number of fleas per rat.
[2] Percent of rats with fleas.

Table 2. *Seasonal changes in abundance of* Leptopsylla segnis *(1944–45)*

	May–June			July–August			September–October			November–December		
	Rats	Mean [1]	Percent [2]	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	4	------	------	27	0	0	15	0	0	6	0.8	50
Immature	8	0	0	31	0	0	29	0	0	26	.8	4
Norway Rats:												
Adults	27	4.0	37	43	.2	2	22	.4	4	12	.4	8
Immature	13	3.8	31	13	0	0	17	.6	6	2	------	------

	December–January–February			March–April			May–June			July–August		
	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	53	1.2	37	52	2.6	75	22	0.4	18	19	0	0
Immature	58	1.5	36	77	2.2	45	63	.4	27	45	0	0
Norway Rats:												
Adults	27	.2	18	60	7.1	63	19	9.0	47	57	0	0
Immature	5	1.2	20	34	9.4	65	16	.6	37	40	0	0

[1] Mean number of fleas per rat.
[2] Percent of rats with fleas.

reaches a peak from May to August and drops to a minimum in the winter. The indices are shown to be higher in July and August 1944, than for the same months in 1945.

Table 2 shows the seasonal changes in abundance for *L. segnis*. The maximum abundance occurs in May and June and the species is practically absent in July and August.

The chicken flea, *E. gallinacea*, was frequently found on rats at all seasons of the year, but no conclusions concerning seasonal abundance can be made from this study as no attempt was made to get monthly samples of rats from poultry houses or chicken coops.

N. fasciatus was found in very small numbers (total of 25) most commonly in March and April. None was found in July and August.

Ctenocephalides sp. was found regularly in suitable places. No conclusions as to seasonal abundance can be drawn from the collections on rats, but judging from the number of complaints to the health department of flea infestations in yards and houses, it seems that cat fleas are most common in May and June.

The human flea, *P. irritans*, is rare on rats. A total of eight was collected, mostly between April and July.

From these observations on fleas, it is clear that the maximum abundance occurs in May and June, but that, because of different peaks of abundance of two species, fleas are common in San Antonio for about 6 months.

For the comparison of abundance of ectoparasites, the rats were classified according to three types of establishments, residences, stores, (cafes, groceries, theaters, drug stores) and mills (grain and peanut

mills) in order to group the rats from similar habitats. The data show no consistent differences in the numbers of fleas on rats from different types of establishments (tables 1 and 2). These studies, however, were not planned to clarify this problem. Especially designed collections of rats from carefully selected establishments are required to determine if differences exist. It appeared true, however, that greater variation from rat to rat in the number of fleas was present in grain mills than elsewhere. Totals of over a hundred fleas on a rat are common in such places. The record of 423 fleas (mostly *L. segnis*) was obtained on a crippled rat caught in a peanut mill.

In addition to seasonal changes in abundance, other aspects may affect the role of fleas in typhus fever. The total fleas, according to species and sex of rat are shown in table 3 which includes all adult rats irrespective of season or place. It is at once clear that the Norway rat in the San Antonio area had more fleas than did the roof rat.

Table 3. *Total fleas on rats—San Antonio*

Species	Sex	Number rats	Arithmetic		Geometric		Percent with fleas
			Mean	Standard error	Mean	Standard error	
Norway	Male	147	11.84	1.46	4.93	0.051	85
	Female	138	9.79	1.19	3.66	.055	69
Roof	Male	95	4.68	.71	2.39	.060	76
	Female	117	2.69	.45	1.43	.045	65

The number of fleas (*X. cheopis*) found on rats caught outside buildings is known to be small. In this investigation few rats were caught outside of buildings, but in two places rats were living out in the grass and feeding in nearby sheds. Rats caught in these places did not have fleas. However, rats caught near the buildings in which they live had as many fleas as rats caught in the buildings.

It is of interest to compare the number of fleas found on male with those found on female rats. Table 3 presents the arithmetic means and their standard errors for all adult Norway rats and roof rats of each sex. In both cases the males had more fleas than did the females. However, the fact that the number of fleas observed are not in a normal distribution, and that the arithmetic mean is very much influenced by the occasional extreme values in these distributions, makes the mean hard to interpret. An alternative centering constant free of this difficulty is the geometric mean (table 3).

For roof rats the difference between the sexes is 2.99 times its standard error, and hence is considered significant. For Norway rats the difference between sexes is only 1.71 times the standard error of the difference, but the fact that the difference is in the same direction

as in roof rats lends support to the idea that male rats have significantly more fleas than do females in this situation.

The sex ratio of the fleas is of possible importance in the knowledge of the life history of the fleas. The ratios for *X. cheopis* and for *L. segnis* are given in table 4. Analysis of the sex ratios in different types of establishments (not presented here) give very similar results. The recent work of Cole (*9*) shows a correlation between the temperature and the sex ratio of fleas (*X. cheopis*) found on rats. On cold days there are more females than males and on warm days there are more males than females. The sexes were equally represented at about 75° F. in several towns in the southern United States. A seasonal calculation for San Antonio confirms this observation (table 5). The percent of female fleas on Norway rats in the warmer seasons was lower than that in the colder seasons. It is interesting to note that apparently the same temperature relation occurs in *L. segnis* which is collected only in the colder months and has a very small proportion of males (tables 4 and 5).

Table 4. *Sex ratio of* Xenopsylla cheopis

	Rats	Mean of male fleas per rat	Mean of female fleas per rat	Total fleas	Percent male fleas
Roof rats:					
Adults	104	1.9	2.5	457	43
Immature	175	2.7	3.3	1,049	43
Norway rats:					
Adults	208	5.7	5.4	2,301	51
Immature	137	5.2	4.2	1,386	51

Sex ratio of Leptopsylla segnis

	Rats	Mean of male fleas per rat	Mean of female fleas per rat	Total fleas	Percent male fleas
Roof rats:					
Adults	54	1.1	2.3	186	34
Immature	71	1.3	2.5	271	34
Norway rats:					
Adults	65	5.8	9.5	991	38
Immature	33	4.6	15.2	393	39

The sex ratios for both species of fleas differ on the two kinds of rats. Roof rats have fewer male fleas than do Norway rats (table 4). The explanation may be that when fleas leave the rats, they gravitate toward the floor and, assuming that male fleas spend less time on rats than do female fleas, then there should be fewer male fleas in the upper parts of buildings. Therefore, roof rats should have a smaller proportion of male fleas than do Norway rats, as is the case. Since *L. segnis* is collected only in the cold months, it is obvious that more female fleas will be found on the rats.

Table 5. *The percentage of females of two species of fleas by seasons*

Months 1944–45	*Xenopsylla cheopis*				*Leptopsylla segnis*			
	Roof rats		Norway rats		Roof rats		Norway rats	
	Fleas	Percent females	Fleas	Percent females	Fleas	Percent females	Fleas	Percent females
May–June	79	66.0	471	43.7	7	----------	139	51.7
July–August	267	54.3	579	46.4	0	----------	1	----------
September–October	80	48.8	126	36.9	0	----------	1	----------
November	18	50.0	4	----------	3	----------	39	71.7
December–January–February	83	53.0	95	63.2	128	69.5	21	66.6
March–April	166	66.3	430	63.6	283	64.0	964	67.8
May–June	302	49.0	438	47.4	37	67.5	41	63.4
July–August	426	51.4	779	46.9	0	----------	0	----------
September–October	62	45.2	556	45.7	8	----------	3	----------

Observations on Mites

The following observations on mites are included in order to add to our knowledge of the ectoparasites of rats. *Liponyssus bacoti*, the tropical rat mite, the most common mite in San Antonio, is at no time as common as are fleas. *Laelaps nuttali* is present in fair abundance; possibly some individuals of *Eulaelaps stabularis* are included in these totals because of misidentification. *Echinolaelaps echidninus* is a rare species. Frequently, the mites may occur in great numbers on a rat. If the rat has any mites, it usually has either very few or very many.

The population changes are grouped in accordance with the seasons (tables 6 and 7). The peak in numbers of *L. bacoti* is in March and April and the minimum occurs in July and August. Some individuals are present in all months of the year. *Laelaps* in contrast, reaches maximum abundance in July and August and a minimum in the winter months, although it is not common at any time. The few *Echinolaelaps echidninus* were found principally in the spring.

The roof rats appear to have more *L. bacoti* than do the Norway rats. In contrast *Laelaps* is more common on Norway rats than on roof rats (tables 6 and 7). There is no consistent difference between the number of mites found on immature and on adult rats.

Observations on Lice

The rat louse, *Polyplax spinulosa*, was frequently present on rats and sometimes in large numbers. When present in numbers, many hundred or even more than a thousand lice could be collected. One Norway rat was gray in color because of the large number of lice and nits. In these investigations an arbitrary total of 50 was recorded when lice were abundant, because the task of collection and counting was tremendous. Hence the data have limited usefulness.

Table 6. *Seasonal changes in abundance of* Liponyssus bacoti *(1944–45)*

	May–June			July–August			September–October			November		
	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	4			27	0.1	7	15	3.9	60	12	5.7	83
Immature	8	0.9	38	32	.3	12	29	1.2	.4	27	3.8	59
Norway rats:												
Adults	22	4.1	32	44	.2	9	22	1.0	45	17	2.0	70
Immature	13	4.0	28	13	.6	39	17	4.2	35	4		

	December–January–February			March–April			May–June			July–August		
	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	49	10.0	80	62	14.6	80	12	18.0	66	18	1.0	28
Immature	59	8.7	75	75	12.9	80	57	3.8	60	43	1.3	44
Norway rats:												
Adults	26	3.3	46	60	10.0	85	19	.7	31	65	.4	9
Immature	7	14.5	71	35	10.9	74	16	2.4	44	32	.1	6

Table 7. *Seasonal changes in abundance of* Laelaps nuttali *(1944–45)*

	May–June			July–August			September–October			November		
	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	4			27	0	0	15	0.1	7	12	0	0
Immature	8	0.1	12	32	.2	6	29	1	10	27	.2	18
Norway rats:												
Adults	22	.2	9	44	.1	7	22	.5	23	17	.5	23
Immature	13	.5	28	13	1.1	15	17	.2	17	4		

	December–January–February			March–April			May–June			July–August		
	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	49	0.6	4	62	0.1	2	12	0	0	18	0.1	6
Immature	59	.1	7	75	.1	8	57	.1	2	43	.1	4
Norway rats:												
Adults	26	.1	11	60	.3	16	19	0	0	65	1.7	37
Immature	7	0	0	35	.3	15	16	.4	19	32	1.7	40

The abundance is recorded according to seasons (table 8). The maximum occurs in the winter season and the minimum in the summer, but lice are present at all seasons. Norway rats have more lice than do roof rats, as shown by the total percentage calculated from the seasonal percentages in order to take account of different numbers of rats in each season.

Interrelations of Rats, Fleas, and Typhus

The differences in seasonal abundance of the various species of ectoparasites are such that there is considerable overlapping and

972999—51——2

at no time does the number of ectoparasites reach the vanishing point. It will be seen that the maximum abundance of fleas begins in May and lasts till August (tables 1 and 2). This increase is 2 months before the maximum number of typhus cases is noted in humans in San Antonio. (Table 9, compiled from the Weekly Morbidity Reports, shows the trend of cases in San Antonio by 4-week periods for 1944.) This lag seems to be more than would be expected if the fleas are equally infected with typhus at all seasons of the year. It seems possible that in the spring the percent of fleas that are infected, as well as the number of fleas, is increasing.

Table 8. *Seasonal changes in abundance of lice*

Season	Species of rat	Rats combed	Percent of rats infested
May–June 1944	Roof	12	8
	Norway	38	71
July–August 1944	Roof	59	3
	Norway	57	10
September–October 1944	Roof	49	14
	Norway	39	31
November 1944	Roof	39	33
	Norway	20	30
December–January–February 1944–45	Norway	113	36
	Roof	33	73
March–April 1945	Roof	140	37
	Norway	94	30
May–June 1945	Roof	90	25
	Norway	35	37
July–August 1945	Roof	63	21
	Norway	89	33
Total	Roof	565	21.8
	Norway	405	39.4

These observations of seasonal abundance are obviously based upon the abundance of ectoparasites on rats which is the important point in enzootic studies. However, the relation between the number of ectoparasites on rats and the number in a building (absolute abundance) needs investigation. It is not known that the seasonal changes herein described represent the changes in absolute abundance in a building. Clarification of this problem would help to explain the persistence of typhus from year to year. Recent work (*9*) refers to this subject and it may be concluded that the increase in fleas on rats in the warm months is in part due to the greater frequency of feeding by the males.

The maintenance of murine typhus fever obviously depends primarily on the frequency of contacts among the rats and fleas. This frequency will depend upon the number of rats and upon the number of fleas on each rat. It is of considerable importance to know the threshold of flea abundance below which typhus cannot maintain itself. Some observations upon this problem have been made in San Antonio by analysis of complement fixation test results and flea indices. The numbers of ectoparasites found on "positive" and

"negative" rats are shown in table 10. It is clear that there is a difference in the numbers for positive and negative rats. This situation occurs in spite of the fact that typhus has occurred in nearly every building with rats.

Table 9. *Typhus cases reported in San Antonio, 1944*

Month	4-week period	Typhus cases	Month	4-week period	Typhus cases
January	1	2	July	8	20
February	2	2	August	9	11
March	3	2	September	10	14
April	4	3	October	11	17
May	5	4	November	12	6
June	6	3	December	13	5
June-July	7	5			

A comparison of the differences in the numbers of fleas on immature positive and immature negative rats with the numbers on the corresponding adults is of interest. By averaging the totals for both sexes, it is found that, in the Norway rat, the positive adults have 1.85 times as many fleas as the negative adults although the positive immature rats have only 1.37 times as many fleas as the negative immature rats. This is to be expected since immature rats may be living in heavily infested buildings but have not lived long enough to have become infected. A similar situation occurs in the roof rats. Studies of this type suffer the deficiency of comparing the numbers of fleas on rats at a time after the rat became infected and are hence only a suggestive relation.

Table 10. *Number of fleas found on "positive" and "negative" rats in San Antonio (1944–45)*

	Positive complement fixation test						Negative complement fixation test					
	Male			Female			Male			Female		
	Rats	Mean[1]	Percent[2]	Rats	Mean	Percent	Rats	Mean	Percent	Rats	Mean	Percent
Roof rats:												
Adults	28	4.4	64	34	4.4	62	37	2.6	73	68	2.2	65
Immature	4	------	------	29	5.7	83	83	3.9	51	84	2.7	57
Norway rats:												
Adults	43	13.3	84	53	10.5	76	57	7.5	74	53	5.2	66
Immature	20	15.3	95	23	9.2	87	37	6.5	84	35	11.4	86

[1] Mean number of fleas per rat.
[2] Percent of rats having fleas.

All surveys of presence of antibodies in rats are biased in favor of positive rats because typhus is more likely to be present where there are many rats and rats are more easily caught in such places. A survey of rats in places where rats are rare would be difficult but would probably show all negative rats.

Summary

Commensal rats were collected in San Antonio, Tex. from May 1944 to September 1945 and examined for ectoparosites.

The rat flea, *Xenopsylla cheopis*, is most abundant on rats from May to August. The mouse flea, *Leptopsylla segnis*, is most abundant in March and April and is almost absent in the summer. The chicken flea, *Echidnophaga gallinacea*, is frequently found and the northern rat flea, *Nosopsyllus fasciatus*, was present in small numbers in the spring. A few cat fleas (*Ctenocephalides* sp.) and human fleas (*Pulex irritans*) were collected. The maximum abundance of fleas occurs in the spring season, but because of overlapping of maxima for various species, fleas are abundant for almost 6 months.

Norway rats have more fleas than have roof rats. For adult rats of both species, the males had more fleas than the females. The number and sex ratios of fleas in stores, mills, and residences were very similar. For both *X. cheopis* and *L. segnis* a high sex ratio in favor of females occurred in the cool seasons.

The mite, *Liponyssus bacoti*, was most abundant in March and April. *Laelaps nuttali* was most abundant in July and August. A few *Echinolaelaps echidninus* were found in May and June.

The rat louse, *Polyplax spinulosa*, was most common from December to February.

Because of the overlapping of months of abundance, ectoparasites are present on rats in all months of the year. The increase in human typhus cases lags about 2 months behind the increase in flea abundance. Rats positive for typhus have more fleas than have rats negative for typhus.

REFERENCES

(*1*) Davis, David E., and Pollard, Morris: The distribution of murine typhus in rats and in humans in San Antonio. Am. J. Trop. Med. **26:** 619–624 (1946).

(*2*) Davis, David E., and Pollard, Morris. Prevalence of typhus complement-fixing antibodies in human serums in San Antonio, Tex. Pub. Health Rep. **61:** 928–931 (1946).

(*3*) Davis, David E.: The control of rat fleas (*Xenopsylla cheopis*) by DDT. Pub. Health Rep. **60:** 485–489 (1945).

(*4*) Davis, David E.: The use of DDT to control murine typhus fever in San Antonio, Tex. Pub. Health Rep. **62:** 449–463 (1947).

(*5*) Davis, David E.: Observations on rats and typhus in San Antonio, Tex. Pub. Health Rep. **63:** 783–790 (1948).

(*6*) Mohr, Carl O.: Results of cooperative State-Federal typhus control programs. CDC Bull. **9:** 1–5 (1950).

(*7*) Hill, Elmer L., and Ingraham, S. C.: A study of murine typhus fever in Coffee County, Alabama. Pub. Health Rep. **62:** 875–881 (1947).

(*8*) Rumreich, A. S., and Koepke, J. A.: Epidemiological significance of seasonal variations in rodent-ectoparasite distribution. Pub. Health Rep. **60:** 1421–1428 (1945).

(*9*) Cole, L. C.: The effect of temperature on the sex ratio of *Xenopsylla cheopis* recovered from live rats. Pub. Health Rep. **60:** 1337–1342 (1945).

Use of Warfarin-Treated Oats as a Plague Suppressive Measure in Hawaii

By Bertram Gross, Robert H. Baker, and David D. Bonnet*

The rodenticidal possibilities of warfarin-treated oats under Hawaiian field conditions were investigated by Doty (*1*) in three tests conducted on the island of Kauai. The results of these field tests showed that warfarin was readily accepted by rodents in this form and that the period of exposure in bait stations should be at least 17 days or until consumption reaches zero. Doty concluded that warfarin had a place in the over-all plantation field program. He suggested that a system using lines of fairly permanent poison-bait stations along gulches and wasteland, where migration is more or less continuous, would be very effective.

Eskey (*2*) pointed out that *Rattus hawaiiensis* undoubtedly plays an important role in maintaining the endemic rural type of rodent plague which is present in Hawaii. The determination of the value of warfarin-treated oats as an additional plague suppressive measure in field areas where large numbers of *R. hawaiiensis* were present became, therefore, a matter of considerable interest and importance to plague control workers in the Territory of Hawaii.

To evaluate this problem in a known endemic plague region in Hawaii, a field experiment was undertaken to find out what effect the continuous application of warfarin-treated oats, over an extended period of time, would have on the rodent population of a limited area.

The area selected for this purpose, field 109, is located in plague work zone 3A, Hamakua, Hawaii. This field (fig. 1) was chosen because reports received from plantation field workers indicated that rats had been causing severe cane damage, and the Bureau of Rodent Control trapping records showed that a high rodent population was present.

Approximately 78 acres of this land are utilized by the Honokaa Sugar Company to cultivate sugar cane. At the time the experiment was initiated, the sections under cultivation were covered with a dense growth of mature cane (variety 32–8560) which was 17 months old and was scheduled to be harvested in a few months. The remainder of the area, approximately 60 acres, was not under cultivation. In these wasteland sections many species of fruit trees, shrubs, and grasses are found, several of which, in addition to sugar cane, may serve at one time or another as a source of rodent food supply.

*Chief, Bureau of Rodent Control, Supervising Rodent Control Inspector, Hamakua, and Medical Entomologist, respectively, Division of Sanitation, Department of Health, Territory of Hawaii.



Figure 1. Field 109, Hamakua District, Hawaii, T. H., selected for rodenticidal experiment.

Many rock piles and a few rock walls are scattered throughout the wasteland sections. The experiment began February 3, 1951, and ended June 8, 1951.

Methods

To obtain a relative measurement of the rodent population, 200 numbered snap traps baited with coconut squares were set at intervals

of 35 to 50 feet throughout the area. These traps were serviced each working day, and records were kept of the rodent catch by individual station. At the end of a 1-week period the traps were removed and portable bait stations were substituted. These stations consisted of empty sardine cans (6½ x 4¼ x 1½ inches) and a numbered, curved covering hood (12 x 13 inches) similar to that described by Doty (*3*).

Unpoisoned oats were not used at any time during this experiment. When the poison stations were first distributed, 4 ounces by weight of warfarin-treated oats [1] were placed in each bait pan. Thereafter, all stations were checked at 3-to 5-day intervals. The presence of fecal pellets, blood stains, and dead rats was noted, and bait consumption was recorded. The bait supply was replenished up to 4 ounces whenever indicated. This poisoning phase of the test was carried on for 91 consecutive days. At the end of this time the poison stations were replaced with snap traps. Snap trapping was continued for 4 weeks to obtain data that could be compared with pretrapping data. Unfortunately post-trapping activities could not be conducted over a longer period of time as field 109 was harvested a few days later.

Results

The total number of rodents retrieved during the pre-poisoning period show that field 109 was heavily infested (table 1). In addition, the trapping data indicate that *R. alexandrinus* was the predominant species of rat present in the area, that large numbers of *R. hawaiiensis* were also present, that *R. norvegicus* and *R. rattus* were less abundant, and that approximately half (47.9 percent) of the rodents found in the area were mice.

When the warfarin-treated oats were exposed, many stations became active within a few days (table 2). Throughout the poisoning

Table 1. *Species composition of trapped rodents*

Species	Pre-poisoning period		Post-poisoning period [1]							
	First week		First week		Second week		Third week		Fourth week	
	Number trapped	Percent	Number trapped	Percent	Number trapped	Percent	Number trapped	Percent	Number trapped	Percent
R. alexandrinus	76	24.28	4	4.26	7	4.93	3	2.78	7	4.14
R. rattus	8	2.56	0	0	0	0	0	0	0	0
R. norvegicus	21	6.71	2	2.13	0	0	5	4.63	4	2.37
R. hawaiiensis	58	18.53	82	87.23	111	78.17	73	67.59	120	71.00
Total Rats	163	52.08	88	93.62	118	83.10	81	75.00	131	77.51
M. musculus	150	47.92	6	6.38	24	16.90	27	25.00	38	22.49
Total Rodents	313	100.00	94	100.00	142	100.00	108	100.00	169	100.00

[1] Warfarin-treated rolled oats exposed for 91 consecutive days.

[1] The poisoned oats consisted of a commercial product, prepared according to the formula of Doty, and contained by weight 0.025 percent warfarin, 11.0 percent mineral oil, 0.25 percent para-nitro-phenol (a mold deterrent), and 88.73 percent rolled oats.

Table 2. *Average daily consumption of warfarin-treated oats*

[200 bait stations]

Interval (days)	Number stations showing some acceptance	Percent stations showing some acceptance	Total bait consumed (gm.)	Average daily consumption (gm.)
4	145	72.5	3,997	999
3	159	79.5	7,002	2,334
3	185	92.5	9,830	3,277
4	162	81.0	6,790	1,698
3	131	65.5	3,352	1,117
4	154	77.0	5,401	1,350
3	135	67.5	4,132	1,377
4	131	65.5	4,061	1,015
3	160	80.0	4,210	1,403
4	135	67.5	4,387	1,097
3	139	69.5	4,097	1,366
3	101	50.5	2,814	938
5	169	84.5	7,258	1,452
3	131	65.5	3,558	1,186
3	151	75.5	3,877	1,292
4	161	80.5	5,705	1,426
3	122	61.0	3,622	1,207
4	149	74.5	5,521	1,380
3	138	69.0	4,897	1,632
4	175	87.5	7,655	1,914
3	146	73.0	4,486	1,495
4	171	85.5	8,115	2,029
3	170	85.0	6,350	2,117
4	154	77.0	7,782	1,946
3	150	75.0	6,081	2,027
4	185	92.5	8,455	2,114

phase, 316 pounds of the bait were consumed, and a high percentage of the stations continued to remain active. During this period, rodent droppings were observed in the bait pans on 549 different occasions. In addition, blood-stained oats were noted 28 times, and 11 rodents, showing signs of having consumed the poisoned grain, were found dead in, or immediately adjacent to, the stations. There was no evidence that any animal except rodents consumed the bait.

The average daily total consumption of the warfarin-treated oats remained at a low level for the first 4 days of exposure (fig. 2). During the following 6 days, consumption rose sharply. The two subsequent recordings, covering a period of 7 days, show that there was a sharp decline in the amount of poison bait eaten. At this time (17 days), the average daily total consumption of bait was 1,117 grams. Thereafter, the consumption curve showed a series of fluctuations with a gradual upward trend, indicating that large numbers of rodents were still present in the test area.

Examination of the results obtained during the post-trapping period shows that a radical change had occurred in the species composition of the rodents present in the area. The first week, there were marked reductions in the numbers of trapped *R. alexandrinus*, *R. rattus*, *R. norvegicus* and *M. musculus*. In contrast, the number of trapped *R. hawaiiensis* increased. Results obtained for the succeeding 3 weeks show that the species composition of the rats in the area remained relatively the same. The number of mice retrieved during this period showed a gradual increase. These data



Figure 2. Consumption of warfarin-treated oats at 200 bait stations, field 109, Hamakua District, Hawaii, T. H., February 9 to May 11, 1951.

indicate clearly that the application of warfarin-treated oats resulted in control of all rodent species present except the native Hawaiian rat, *R. hawaiiensis*.

Discussion

Results of cage tests undertaken in Honolulu approximately 6 weeks after the Hamakua field experiment had begun, demonstrated that *R. hawaiiensis* accept warfarin-treated oats and die from the effects (see article following). The poison mix utilized for these cage tests was identical with that offered in the field experiment. The cage-test animals, however, did not have an extensive choice of food.

All species of rodents which were present in the field test area had a multiple choice of foods. After the poison bait was distributed, undoubtedly many of these rodents continued to feed to some extent on the various types of food normally available to them. As effective control of *R. alexandrinus*, *R. rattus*, *R. norvegicus*, and *M. musculus* was achieved through the use of warfarin-treated oats, it is apparent that most of these rodents also found the poison bait attractive and ingested quantities which eventually killed them.

The daily consumption of poison bait continued to remain at a high level throughout the poisoning period, indicating that *R. hawaiiensis* was also feeding on the warfarin mix. This conclusion is supported by the fact that 59.8 percent of the *R. hawaiiensis* rats trapped during the first week of the post-poisoning period showed evidence of hemorrhage on autopsy. Since the Hawaiian rat was not controlled effectively, it is not unlikely that this species fed sporadically on the warfarin-treated oats but did not consistently ingest sufficient quantities to produce death.

It is quite possible that the failure of the warfarin preparation to

control *R. hawaiiensis* is related to the question of food preferences. In the field, *R. hawaiiensis* may be more selective in its choice of foods than the other species of rodents. In addition, this species may consistently prefer a greater variety of foods. These speculations indicate the desirability of conducting further investigations to determine the food preferences of *R. hawaiiensis*.

Another possibility is that the *R. hawaiiensis* population was not controlled because of extensive migrations from adjacent areas. However, if one assumes this, it is difficult to explain why the other species of rodents were so drastically reduced. Perhaps the increase in the number of *R. hawaiiensis* was partially related to the radical decrease in the number of the *R. alexandrinus*, *R. rattus* and *R. norvegicus*. Since these species are larger and undoubtedly compete with the Hawaiian rat, it is not improbable that a reduction in their numbers would result in *R. hawaiiensis* being subjected to less biological pressure, thereby giving this species a much greater chance of survival. This would apply to both *R. hawaiiensis* present in the area and to those migrating into it. The younger rats of this species would be benefited particularly by such conditions.

R. hawaiiensis is primarily a field rodent and prefers to live in rock piles or in thickly vegetated areas. It is timid and rarely infests human habitations. The other species of rodents are also found in the fields. However, they are more aggressive and occasionally infest buildings where they and their fleas come into intimate contact with people. In the communities, therefore, *R. alexandrinus*, *R. rattus*, *R. norvegicus*, and *M. musculus* are more directly implicated in the transmission of plague to humans than is *R. hawaiiensis*.

As previously noted, warfarin-treated oats effectively controlled all species of rodents in the test area except *R. hawaiiensis*. It is probable, therefore, that the extensive application of this rodenticide in field areas and in communities might have considerable value as a means of reducing the plague potential for the entire endemic region. However, large numbers of *R. hawaiiensis* are found in all areas of the plague district and plague infection is detected more often in this species than in any other rodent present. Thus, *R. hawaiiensis* and its ectoparasites would continue to be the reservoir of plague infection in the cane fields and wastelands.

The important role played by *R. hawaiiensis* in the epidemiology of plague in Hawaii focuses attention on the necessity of conducting additional laboratory and large-scale field tests to determine a suitable warfarin bait which will effectively control this species.

Summary

1. The purpose of this experiment was to determine the value of warfarin-treated oats as an additional plague suppressive measure

in field areas in Hawaii where large numbers of *Rattus hawaiiensis* were present.

2. The test was conducted in a sugar cane field located in the endemic plague region of the Hamakua District, island of Hawaii, T. H.

3. Trapping results obtained before warfarin-treated oats were exposed show that this field was heavily infested with rodents. The results indicated that *R. alexandrinus* was the predominant species of rat present in the area, that large numbers of *R. hawaiiensis* were also present, that *R. norvegicus* and *R. rattus* were less abundant and that approximately half (47.9 percent) of the rodents found in the area were *M. musculus*.

4. Warfarin-treated oats were exposed continuously at 200 bait stations for a period of 91 days. During this time 316 pounds of bait were consumed, and a high percentage of the stations remained consistently active, indicating that large numbers of rodents were still present in the test area.

5. Trapping results obtained immediately after poisoning activities showed that a radical change had occurred in the species composition of the rodents present in the area. There were marked reductions in the numbers of *R. alexandrinus*, *R. rattus*, *R. norvegicus*, and *M. musculus*. In contrast, the number of trapped *R. hawaiiensis* increased.

6. The data indicate clearly that the application of warfarin-treated oats resulted in control of all rodent species present except the native Hawaiian rat, *R. hawaiiensis*.

7. The extensive application of warfarin-treated oats in field areas and communities to control *R. alexandrinus*, *R. rattus*, *R. norvegicus*, and *M. musculus* might have considerable value as a means of reducing the plague potential for the entire endemic region. However, *R. hawaiiensis* and its ectoparasites would continue to be the reservoir of plague infection in the fields. It is indicated, therefore, that a more suitable warfarin bait be found which will effectively control this species.

REFERENCES

(*1*) Doty, R. E.: Warfarin (Compound 42) A promising new rodenticide for cane fields. Hawaiian Planters' Rec. **54**: 1–21 (1950).
(*2*) Eskey, C. R.: Epidemiological Study of Plague in the Hawaiian Islands. Public Health Bulletin No. 213, pp. 1–70 (1934).
(*3*) Doty, R. E.: Rat control on Hawaiian sugar cane plantations. Hawaiian Planters' Rec. **49**: 74–239 (1945).

Cage Tests With Warfarin on the Hawaiian Rat, *Rattus hawaiiensis* Stone, and the House Mouse, *Mus musculus* Linn., in Hawaii

By David D. Bonnet, Edward S. C. Mau, and Bertram Gross*

A previous report by Doty (*1*) has given the results obtained with warfarin-treated rolled oats as a rodenticide in cage tests in Hawaii on *Rattus norvegicus* (Erxleben), *Rattus alexandrinus* (Geoffroy), and *Rattus rattus* Linn. This paper reports a similar series of tests undertaken in Honolulu, T. H. on the Hawaiian rat, *Rattus hawaiiensis* Stone, and the house mouse, *Mus musculus* Linn. Data on the effectiveness of warfarin-treated rolled oats against these two additional species were particularly desired because an extensive field experiment in which this poison bait was used was already in progress in a rural plague area, where large numbers of both species were known to be present (see preceding report in this issue).

Rats and mice were captured during March and April 1951 near Honolulu, in live cage traps baited with squares of coconut meat. They were transferred to individual cages and provided with unlimited drinking water and pieces of coconut. Each cage contained nesting material and a suitable shelter. A record was made of the species, sex, and weight of each rodent.

Tared petri dishes containing weighed quantities of rolled oats were placed in the cages each day. These dishes and their contents were reweighed on each subsequent day and the amount consumed was determined by difference. Unpoisoned rolled oats were presented for approximately seven consecutive days, after which poisoned oats were substituted. Preliminary tests had demonstrated that a pretest period of 1 week was more than sufficient for the caged wild rodents to become accustomed to laboratory life and to consume regular amounts of food. The poisoned oats consisted of a commercial product, prepared according to the formula of Doty, and contained by weight 0.025 percent warfarin, 11.0 percent mineral oil, 0.25 percent para-nitro-phenol (a mold deterrent), and 88.73 percent rolled oats.

---

*Medical Entomologist, former Supervising Rodent Control Inspector, Oahu, and Chief, Bureau of Rodent Control, respectively, Division of Sanitation, Department of Health, Territory of Hawaii.

Note: 3 (alpha-phenyl-beta-acetyl ethyl)-4-hydroxycoumarin, or warfarin, is a chemical discovered, by Dr. K. P. Link and associates of the Wisconsin Alumni Research Foundation, to have lethal anti-coagulant properties in mammals.

Results

A summary of the results obtained is presented in the table. There was a difference in the quantity of food eaten by the male and female rats, but when calculated on a body weight basis, the difference was not significant. The poisoned oats were not quite as acceptable as unpoisoned oats, since the daily food consumption dropped significantly on the first 2 days after presentation of the poison. After the second day there was an even further drop in consumption (fig. 1), probably indicating that the poison was beginning to take effect. The increased daily consumption as shown in figure 1 after the sixth day is the result of consumption by those animals which lived beyond the mean day of death (7.9 days). The animals in this group, constituting 50 percent of the population, ate a smaller quantity of warfarin oats (4.9 gm.) during the first 2 days of the experiment, and were still consuming small quantities of poisoned bait at the same time that early mortalities were eliminating a portion of the population. Therefore, the food consumption, as percent of body weight, is warped and indicates an artificial increase which is not significant. This avoidance of the poisoned bait by some individuals may account in part for the longer survival of this group.

The mean day of death (7.9) is somewhat greater in *R. hawaiiensis* than that reported by Doty for the other species of rodents found in Hawaii (*R. norvegicus*, 5.4 days; *R. rattus*, 5.8 days; and *R. alexandrinus*, 6.8 days). This would appear to indicate a greater resistance of *R. hawaiiensis* to the effects of the poison. However,



Figure 1. Consumption of warfarin-treated oats by *R. hawaiiensis*.

the amount of poisoned oats ingested by this species in percentage of body weight was 38.9±1.3 percent. This quantity is not significantly different from the amounts ingested by *R. alexandrinus* in Doty's tests. None of the animals in this series refused to accept the warfarin-poisoned oats, and all died from the effects.

Results of cage tests with warfarin, in Hawaii, March and April 1951

Item	*Rattus hawaiiensis*	*Mus musculus*
Number	42	15.
Total body weight	2,313.8 gm.	198.2 gm.
Mean body weight	55.1±1.2 gm.	13.2±1.0 gm.
Mean daily consumption of unpoisoned rolled oats	7.1±0.2 gm.	2.3±0.3 gm.
Mean daily consumption as percent of body weight	12.9±0.8 percent.	17.4±3.0 percent.
Total warfarin-oats eaten	899.0 gm.	143.0 gm.
Mean daily consumption of warfarin-poisoned oats on 1st and 2d day of presentation.	5.6±0.2 gm.	1.4±0.4 gm.
Total warfarin-oats as percent of total body weight	38.9±1.3 percent.	72.2±6.5 percent.
Mean day of death	7.9±0.5 days.	8.6±1.0 days.

The earliest that death occurred was 4 days after the initial presentation of poison. The maximum survival was 20 days. By the eleventh day, 90 percent of the experimental animals had succumbed to the effects of warfarin.

Autopsies were performed on all animals after death, and hemorrhages were found in various organs of the body, including the heart,



Figure 2. Consumption of warfarin treated oats by *M. musculus*.

lungs, mesenteries, brain, skin, muscles, and genitourinary systems. In most instances, more than one system was involved. It is noteworthy, however, that 69 percent of the rats showed extensive hemorrhages in the pectoral-cervical region. Hayes and Gaines (*2*) have stated that for the white rat the perioesophageal and cervical bleeding was possibly the result of mild experimental trauma due to stomach tube application of the poison. Our results indicate that in *R. hawaiiensis* cervical involvement occurs naturally and with high frequency.

As would be expected in a smaller animal, the food consumption of mice in terms of percentage of body weight, was greater than that found in the rats (fig. 2). The mean day of death for *Mus musculus* was 8.6 days, which is significantly greater than that found for *Rattus hawaiiensis* or for the three species tested by Doty. The ingestion of warfarin by mice continued for a longer period and resulted in a high consumption of poisoned rolled oats (72.2 ± 6.5 percent) when computed in terms of percentage of body weight. Ninety percent of the mice were dead as a result of the poison by the end of the thirteenth day. The earliest death occurred on the fourth day, and the maximum survival of any individual mouse was 17 days.

REFERENCES

(*1*) Doty, R. E.: Warfarin (Compound 42) A promising new rodenticide for cane fields. Hawaiian Planters' Rec. **54:** 1–21 (1950).

(*2*) Hayes, Wayland J., Jr., and Gaines, Thomas B.: Control of Norway rats with residual rodenticide warfarin. Pub. Health Rep. **65:** 1537–1555. (1950).

Incidence of Disease

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

Announcement

A New *Morbidity and Mortality Weekly Report*

The new monthly PUBLIC HEALTH REPORTS (see back cover) will publish from time to time, as appropriate, reports, tabulations, and articles dealing with morbidity statistics, both domestic and foreign. The present weekly "Incidence of Disease" section, however, will be discontinued as of December 31, 1951.

Current provisional morbidity data on notifiable diseases for the United States now appear in summary form and in tabulations by States and cities in the *Weekly Morbidity Report* issued by the National Office of Vital Statistics of the Public Health Service. Beginning on January 11, 1952, the Public Health Service through the National Office of Vital Statistics will issue a *Morbidity and Mortality Weekly Report* presenting these morbidity data as well as certain mortality data for selected cities.

Libraries and agencies that have depended upon PUBLIC HEALTH REPORTS for current morbidity statistics for the United States may continue to receive the same data by writing to the National Office of Vital Statistics, Public Health Service, Washington 25, D. C., requesting that they be placed on the mailing list for the new *Morbidity and Mortality Weekly Report.* Individuals who wish to be placed on the mailing list should indicate how and to what extent they will make use of this publication.

Since the *Weekly Epidemiological Record* and other publications of the World Health Organization, Geneva, Switzerland, contain morbidity data for foreign countries, tabulations of notifiable diseases occurring outside the United States and its Territories will not appear regularly in National Office of Vital Statistics publications.

UNITED STATES

Reports from States for Week Ended December 8, 1951

The incidence of measles declined slightly for the current week as compared with the previous week but was about 66 percent higher than for the same week in 1950. The greatest concentration of cases continues to be in the northeastern section of the country.

The number of cases of scarlet fever (1,530) for the current week is about 30 percent higher than for the same week last year. The 5-year median is 1,837.

A slight decrease in poliomyelitis cases was reported. The cumulative total for the calendar year is now 27,975 as compared with

32,473 for the same period in 1950. The cumulative total since the seasonal low week is 26,757 as compared with 31,342 in 1950.

Diphtheria incidence was lower for the current week than for the same week last year. The disease continues to be concentrated in the southern States.

Two cases were erroneously reported as smallpox in Nebraska last week because of a clerical error. They were cases of chickenpox.

Nine cases of rabies in animals were erroneously reported for New Jersey for the week ended November 24 because of an error in transmitting the report.

One case of leprosy was reported by Minnesota.

Only eight cases of malaria in civilians were reported, two in New York and six in Texas. The number from military establishments was also smaller as compared with previous weeks.

Epidemiological Reports

Gastroenteritis

Dr. J. P. Ward, Arizona Director of Public Health, has reported an outbreak of gastroenteritis consisting of 40 cases which occurred at a church social. All food stuffs served at the meal were examined bacteriologically and *Staphylococcus* was isolated from a sample of cole slaw. Mayonnaise was suspected of containing the organism and improper refrigeration was thought to be a contributing factor.

Dr. H. W. Stevens, Massachusetts district health officer, has reported an outbreak of gastroenteritis caused by *Salmonella montevideo.* Nine cases occurred after an incubation period of 18 to 24 hours following the eating of chocolate eclairs. The illness lasted from 1 to 2 weeks. Because of a marked delay in reporting the cases, investigation of the bakery where the eclairs were prepared, and of its personnel, did not reveal the source of infection.

S. V. Dugan, Kentucky Department of Health, has reported an outbreak of gastroenteritis in a public school which presumably followed the eating of wieners in the cafeteria. About 45 to 50 pupils out of a total of 98 who ate in the cafeteria became ill after an incubation period of 3 to 10 hours. A sample of uncooked wieners, when examined in the laboratory, showed the presence of a weakly hemolytic alpha prime type *Streptococcus.* Fifty grams of the sample fed to a kitten caused marked diarrhea 24 hours later.

Reports have been received that construction workers and other personnel on a project in Nevada were affected twice within 2 weeks by outbreaks of gastroenteritis, presumably food-borne. Sixty persons were affected in the first and 100 or more in the second instance. Some article of food in box lunches, which included meat sandwiches, is suspected of being the vehicle of infection.

Rabies in Animals

Rabies has been reported in a number of species of animals over the past few months in South Dakota. Prior to 1951, no cases in animals had been observed for about 13 years. Since the disease appeared in April 1951, laboratory confirmation has been obtained on the following species of animals: dogs, cats, skunks, muskrats, squirrels, rats, and mice. The infection has been found more frequently in skunks than in dogs or other domestic animals.

Psittacosis

Dr. Albert Milzer, Collaborating Laboratory of the Influenza Study Program, has reported a complement fixation titer of 1 to 64 in acute phase serum, and 1 to 256 in convalescent serum for the psittacosis-pneumonitis group in a woman who owns a pet shop in Chicago. The patient who sells parrots and parakeets had clinical findings of an atypical pneumonia and responded successfully to aureomycin therapy.

Comparative Data for Cases of Specified Reportable Diseases: United States

[Numbers after diseases are International List numbers, 1948 revision]

Disease	Total for week ended—		5-year median 1946–50	Seasonal low week	Cumulative total since seasonal low week		5-year median 1945–46 through 1949–50	Cumulative total for calendar year—		5-year median 1946–50
	Dec. 8, 1951	Dec. 9, 1950			1950–51	1949–50		1951	1950	
Anthrax (062)	------	1	1	(1)	(1)	(1)	(1)	57	43	48
Diphtheria (055)	90	132	220	27th	1,910	2,573	4,518	3,918	5,701	9,129
Encephalitis, acute infectious (082)	26	20	11	(1)	(1)	(1)	(1)	999	953	608
Influenza (480–483)	1,695	1,210	1,210	30th	8,045	11,097	11,097	124,100	149,864	139,295
Measles (085)	6,059	3,008	3,008	35th	31,341	18,254	18,238	500,252	306,425	584,626
Meningitis, meningococcal (057.0)	76	80	69	37th	804	741	709	3,865	3,540	3,253
Pneumonia (490–493)	1,089	1,551	(2)	(1)	(1)	(1)	(1)	56,221	76,282	(2)
Poliomyelitis, acute (080)	378	480	322	11th	[3] 26,763	31,342	26,666	[3] 27,975	32,473	27,016
Rocky Mountain spotted fever (104)	------	1	1	(1)	(1)	(1)	(1)	328	452	558
Scarlet fever (050) [4]	1,530	1,194	1,837	32d	11,662	11,673	15,898	65,048	51,843	72,259
Smallpox (084)	------	------	1	35th	[5] 1	11	11	[5] 12	37	53
Tularemia (059)	20	16	22	(1)	(1)	(1)	(1)	613	841	962
Typhoid and paratyphoid fever (040,041) [6]	53	54	51	11th	2,553	2,798	3,238	2,988	3,307	3,723
Whooping cough (056)	1,502	1,967	2,227	39th	11,286	16,964	16,964	65,061	114,159	93,755

[1] Not computed. [2] Data not available. [3] Addition: Kentucky, 6 cases, delayed reports—not allocated. [4] Including cases reported as streptococcal sore throat. [5] Deduction: Nebraska, week ended Dec. 1, 2 cases. [6] Including cases reported as salmonellosis.

Reported Cases of Selected Communicable Diseases: United States, Week Ended Dec. 8, 1951

[Numbers under diseases are International List numbers, 1948 revision]

Area	Diphtheria (055)	Encephalitis, infectious (082)	Influenza (480–483)	Measles (085)	Meningitis, meningococcal (057.0)	Pneumonia (490–493)	Poliomyelitis (080)
United States	**90**	**26**	**1,695**	**6,059**	**76**	**1,089**	**378**
New England		**1**	**4**	**1,094**	**5**	**44**	**5**
Maine			2	129		9	1
New Hampshire				46	1	4	
Vermont				137			
Massachusetts		1		439			3
Rhode Island				91	3	1	
Connecticut			2	252	1	30	1
Middle Atlantic	**6**	**4**	**1**	**2,394**	**19**	**102**	**19**
New York	3	4	(1)	1,368	10		9
New Jersey			1	294		40	
Pennsylvania	3			732	9	62	10
East North Central	**19**	**9**	**5**	**1,035**	**14**	**94**	**46**
Ohio	9			230	5		11
Indiana	8	2		33			6
Illinois	1	3	3	306	3	78	6
Michigan	1	4	2	364	6	16	16
Wisconsin				102			7
West North Central	**7**	**1**	**7**	**129**	**5**	**212**	**36**
Minnesota			1	47	1	13	13
Iowa				7	1		2
Missouri	6		1	5	1		6
North Dakota		1	5	35		198	
South Dakota				9	1		1
Nebraska				16			4
Kansas	1			10	1	1	10
South Atlantic	**29**	**3**	**75**	**508**	**8**	**123**	**27**
Delaware				1			
Maryland	1		1	231		33	3
District of Columbia				31		17	1
Virginia	3	2		65	3	40	4
West Virginia	4	1		44	1		3
North Carolina	3			7	4		4
South Carolina	1		44	14		4	
Georgia	13		30	91		29	5
Florida	4			24			7
East South Central	**17**	**1**	**14**	**243**	**10**	**53**	**40**
Kentucky	2		2	168	3	2	5
Tennessee	1	1		44	5		5
Alabama	12			17	1	37	8
Mississippi	2		12	14	1	14	22
West South Central	**8**	**1**	**203**	**81**	**4**	**287**	**70**
Arkansas			97	15		31	13
Louisiana			1	1		31	7
Oklahoma	1		105	6	2	9	2
Texas	7	1		59	2	216	57
Mountain	**1**		**1,357**	**283**	**3**	**85**	**35**
Montana			25	41	1		2
Idaho	1			5			6
Wyoming				1		3	4
Colorado			1,117	24		20	5
New Mexico				10	1	1	1
Arizona			215	145		61	2
Utah				57	1		13
Nevada							2
Pacific	**3**	**6**	**29**	**292**	**8**	**89**	**91**
Washington	2		4	103	1	2	10
Oregon			13	14	2	39	6
California	1	6	12	175	5	48	75
Alaska		1					
Hawaii			32	784			

[1] New York City only.

Reported Cases of Selected Communicable Diseases: United States, Week Ended Dec. 8, 1951—Continued

[Numbers under diseases are International List numbers, 1948 revision]

Area	Rocky Mountain spotted fever (104)	Scarlet fever [1] (050)	Smallpox (084)	Tularemia (059)	Typhoid and paratyphoid fever [2] (040, 041)	Whooping cough (056)	Rabies in animals
United States		**1,530**		**20**	**53**	**1,502**	**129**
New England		**85**			**4**	**200**	
Maine		4				33	
New Hampshire		9				12	
Vermont		2				41	
Massachusetts		48			3	85	
Rhode Island		1				14	
Connecticut		21			1	15	
Middle Atlantic		**258**		**1**	**4**	**216**	**7**
New York		173			3	100	3
New Jersey		19				40	
Pennsylvania		66		1	1	76	4
East North Central		**528**		**6**	**5**	**281**	**10**
Ohio		151			1	100	4
Indiana		64		1	1	13	1
Illinois		99		2	1	29	2
Michigan		171		3	1	81	1
Wisconsin		43			1	58	2
West North Central		**77**		**2**		**85**	**19**
Minnesota		18		2		1	10
Iowa		5				6	4
Missouri		23				26	5
North Dakota		1				4	
South Dakota		3				2	
Nebraska		4				24	
Kansas		23				22	
South Atlantic		**179**		**2**	**7**	**145**	**28**
Delaware							
Maryland		32				9	
District of Columbia		13			1		
Virginia		23		1	2	24	3
West Virginia		17				67	2
North Carolina		69			1	14	12
South Carolina		2					3
Georgia		11		1	3	13	8
Florida		12				18	
East South Central		**49**		**3**	**2**	**109**	**29**
Kentucky		23		3		45	4
Tennessee		25				28	11
Alabama					1	32	6
Mississippi		1			1	4	8
West South Central		**49**		**5**	**9**	**334**	**32**
Arkansas		6		4	1	11	2
Louisiana		4				3	
Oklahoma		9		1	1	11	2
Texas		30			7	309	28
Mountain		**50**		**1**	**3**	**40**	
Montana		10				4	
Idaho		8				11	
Wyoming		1		1		2	
Colorado		7			1	17	
New Mexico		3			1	1	
Arizona		4			1	4	
Utah		12				1	
Nevada		5					
Pacific		**255**			**19**	**92**	**4**
Washington		20			1		
Oregon		29				8	
California		206			18	84	4
Alaska		1					
Hawaii					3		

[1] Including cases reported as streptococcal sore throat. [2] Including cases reported as salmonellosis.

FOREIGN REPORTS

CANADA

Reported Cases of Certain Diseases—Week Ended Nov. 24, 1951

Disease	Total	Newfoundland	Prince Edward Island	Nova Scotia	New Brunswick	Quebec	Ontario	Manitoba	Saskatchewan	Alberta	British Columbia
Brucellosis	2					2					
Chickenpox	1,228	6		33	4	189	508	56	95	185	152
Diphtheria	12					11				1	
Dysentery:											
Amebic	3					3					
Bacillary	3						3				
Encephalitis, infectious	1								1		
German measles	167			82		22	23		12	13	15
Influenza	16			12			2	1	1		
Measles	793	14			6	154	97	20	2	358	142
Meningitis, meningococcal	10					4			1	1	4
Mumps	718	1			2	130	388	38	47	60	52
Poliomyelitis	20			2		8	3		5	1	1
Scarlet fever	385				2	122	31	23	44	44	119
Tuberculosis (all forms)	271	33		10	2	123	23	35	10	12	23
Typhoid and paratyphoid fever	5					5					
Venereal diseases:											
Gonorrhea	301	6		3	3	74	57	26	33	41	58
Syphilis	68	5		5	1	36	10	2	1	1	7
Primary	6					3	1				2
Secondary	5					3	1			1	
Other	57	5		5	1	30	8	2	1		5
Other forms	1							1			
Whooping cough	257					116	60	16	15	32	18

MADAGASCAR

Reported Cases of Certain Diseases and Deaths—September 1951

Disease	Aliens		Native	
	Cases	Deaths	Cases	Deaths
Beriberi			1	
Bilharziasis			86	
Dysentery:				
Amebic	2		153	
Bacillary	2		21	
Diphtheria	1		9	
Erysipelas			6	
Influenza	2		3,388	13
Leprosy			24	
Malaria	103	2	24,824	113
Measles	2		229	4
Mumps			100	
Meningitis, meningococcal			6	3
Plague			25	20
Pneumonia (all forms)	4	1	691	70
Puerperal infection			8	1
Tuberculosis, respiratory	8	1	109	15
Typhoid fever			3	
Whooping cough	7		293	

REPORTS OF CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER RECEIVED DURING THE CURRENT WEEK

The following reports include only items of unusual incidence or of special interest and the occurrence of these diseases, except yellow fever, in localities which had not recently reported cases. All reports of yellow fever are published currently.

Cholera

India. During the week ended December 1, cholera was reported as follows: Calcutta, 77 cases; and Madras, 8.

Pakistan. For the week ended December 1, 20 cases of cholera were reported in Dacca.

Plague

Madagascar. There were more cases (25) of plague reported during the period November 11–20 than for any previous 10-day period this year. Only 9 cases were reported for the period November 1–10.

Union of South Africa. One case of septicemic plague was reported during the week ended November 24, in the Bothaville District, Orange Free State.

Smallpox

Algeria. One case of smallpox was reported during the period October 21–31. This is the first case since August 21.

Burma. Smallpox was reported for the week ended December 1, as follows: Mergui, 27 cases; Moulmein, 2; and Rangoon, 1.

India. During the week ended December 1, smallpox was reported in ports of India as follows: Madras, 8 cases; Calcutta, 5; Bombay, 2; and Cawnpore, 1.

Indochina. The incidence of smallpox in Hanoi, Viet Nam, has increased from 33 cases reported for the week ended November 24, to 208 for the following week.

Yellow Fever

Gold Coast. The three cases of yellow fever reported in Suhum for the period August 2–9, were confirmed on November 30.

✦ ✦ ✦

The printing of this publication has been approved by the Director of the Bureau of the Budget (August 10, 1949).

The PUBLIC HEALTH REPORTS, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the Public Health Service through the Division of Public Health Methods, pursuant to the following authority of law: United States Code, title 42, sections 241, 245, 247; title 44, section 220.

It contains (1) current information regarding the incidence and geographic distribution of communicable diseases in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The PUBLIC HEALTH REPORTS is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work.

Requests for and communications regarding the PUBLIC HEALTH REPORTS and reprints should be addressed to the Surgeon General, Public Health Service, Washington 25, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington 25, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.

✦ ✦ ✦

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON D. C. : 1951

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C.
Price 10 cents. Subscription price—per year $4.25, domestic; $5.00 foreign.

—Announcement—

PUBLIC HEALTH REPORTS
To be Published Monthly Next Year

Beginning in January 1952 PUBLIC HEALTH REPORTS will be published monthly. Two other technical publications of the Public Health Service—the Journal of Venereal Disease Information and the CDC Bulletin—are being merged with it.

The merger has come about as the result of an extended study of the Public Health Service's publishing activities. The new journal is designed to carry out more efficiently and economically the Public Health Service's responsibilities for disseminating scientific information on matters of health.

The new PUBLIC HEALTH REPORTS will be concerned with the technical and professional aspects of public health practice, with problems of health administration, and with research in these fields. It will include substantially the types of material that have been appearing in the three present journals and the monthly Tuberculosis Control Issue of PUBLIC HEALTH REPORTS. There will be more emphasis, however, on administrative practice, program development, and applied research aspects and less emphasis on bench research and clinical material not rather directly related to public health practice.

As has been true with each of the present journals, the pages of the new PUBLIC HEALTH REPORTS will be open, on merit, to any responsible author. General editorial guidance will be provided by a Board of Editors composed of individuals of recognized competence and professional stature drawn from both within and outside the Federal service.

Every effort will be made to furnish the new PUBLIC HEALTH REPORTS to representative organizations in public health and its related fields. Most of the readers on the mailing list are affiliated with organizations to which free distribution can legally be made. They will continue to receive the new journal either directly or through these organizations.

The first issue of the new PUBLIC HEALTH REPORTS and an application form for a free subscription will be sent to readers now on the free lists for technical periodicals. After they have had an opportunity to review the new PUBLIC HEALTH REPORTS and to consider its usefulness in their work, readers who wish to remain on the lists and receive the new periodical regularly should complete the application and send it in as soon as possible. This will allow time to revise the lists for mailing the second and subsequent issues.

The new PUBLIC HEALTH REPORTS will also be available by paid subscription from the Superintendent of Documents, Government Printing Office.

PUBLIC LIBRARY
FEB 5 1952
DETROIT

SOCIAL SCIEN

Public Health Reports

Issued Weekly by the
PUBLIC HEALTH SERVICE

Index

Volume 66—Part II

Nos. 27–52

July–December 1951

FEDERAL SECURITY AGENCY

PUBLIC HEALTH SERVICE

FEDERAL SECURITY AGENCY
Oscar R. Ewing, Administrator

PUBLIC HEALTH SERVICE
Leonard A. Scheele, Surgeon General

Division of Public Health Methods
G. St. J. Perrott, Chief of Division

The printing of this publication has been approved by the Director of the Bureau of the Budget (August 10, 1949).

The PUBLIC HEALTH REPORTS, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the Public Health Service through the Division of Public Health Methods, pursuant to the following authority of law: United States Code, title 42, sections 241, 245, 247; title 44, section 220.

It contains (1) current information regarding the incidence and geographic distribution of communicable disease in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The PUBLIC HEALTH REPORTS is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work.

Requests for and communications regarding the PUBLIC HEALTH REPORTS and reprints should be addressed to the Surgeon General, Public Health Service, Washington 25, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington 25, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON : 1952

For sale by the Superintendent of Documents, U. S. Government Printing Office Washington 25, D. C. Index, 10 cents. Subscription price, $4.25 a year, 75 cents additional for foreign mailing; single copies vary in price.

Public Health Reports Index

Subject Index

Key to Dates and Pages

No.	Date of issue	Pages	No.	Date of issue	Pages
27	July 6	851– 882	40	Oct. 5	1263–1302
28	July 13	883– 910	41	Oct 12	1303–1338
29	July 20	911– 950	42	Oct. 19	1339–1378
30	July 27	951– 986	43	Oct. 26	1379–1414
31	Aug. 3	987–1017	44	Nov. 2	1415–1449
32	Aug. 10	1019–1046	45	Nov. 9	1451–1486
33	Aug. 17	1047–1074	46	Nov. 16	1487–1522
34	Aug. 24	1075–1106	47	Nov. 23	1523–1558
35	Aug. 31	1107-1138	48	Nov. 30	1559–1594
36	Sept. 7	1139–1170	49	Dec. 7	1595–1648
37	Sept. 14	1171–1194	50	Dec. 14	1649–1684
38	Sept. 21	1195–1226	51	Dec. 21	1685–1716
39	Sept. 28	1227–1262	52	Dec. 28	1717–1744

A

Page

Absenteeism, industrial sickness 1550
Accident prevention, home 1461
Age incidence of specific causes of illness found in monthly canvasses of families—Sample of the Eastern Health District of Baltimore, 1938–43 [Collins, Phillips, and Oliver] 1227
Agglutination-inhibition test for virus influenza, preparation of dried antigen and antiserum for 1195
Aging, establishment of a new bulletin 910
Alaska public health problems 911, 912, 917, 922, 928, 934, 939, 941
Alaska sewage and waste disposal projects 922
Aleuts, Alaskan, public health problems of 912
Allergy, tuberculin, following BCG vaccination 1415
Altitudes and tuberculin sensitivity 1427
American Public Health Association: Southern Branch conference report 1107
Antigen and antiserum, dried, for the agglutination-inhibition test for virus influenza 1195
Antigens, changes induced in the flagellar, of *Salmonella rostock* and *Salmonella california* 1694
Arctic Health Research Center facilities and opportunities for research 941
Arteriosclerotic coronary artery disease: *See* Occlusion, coronary.
Atomic warfare and the dentist 1175

B

Page

Baltimore, Eastern Health District of, sample of white families canvassed at monthly intervals, 1938–43 ... 1227, 1649
BCG vaccination:
In Poland ... 873
On a world wide scale ... 1158
Research contributions of, programs ... 1427
Tuberculin allergy following ... 1415
Biotic interrelationships of helminth parasitism [Rausch] ... 928
Births, premature, and neonatal mortality statistics ... 1038
Bovine mammary gland Q fever infection ... 1685
Brucella, a yolk sac technique for the routine isolation of. Injection of clotted blood specimens into embryonating eggs with recovery of all three species [Gay and Damon] ... 1204
Brucella melitensis, viability of, in naturally infected cured hams [Hutchings, McCullough, Donham, Eisele, and Bunnell] ... 1402
Bureau of State Services' new chief ... 1676

C

Cage tests with warfarin on the Hawaiian rat, *Rattus hawaiiensis* Stone, and the house mouse, *Mus musculus* Linn., in Hawaii [Bonnet, Mau, and Gross] ... 1734
Calves, effect on, of oral administration of molluscacides ... 1313
Cancer Investigation Center, Hot Springs National Park. Experience with a streamlined examination [Koplin] ... 1339
Caries reduction: *See* Dental caries
Case reporting, tuberculosis ... 1291
Case reports: Coronary occlusion with myocardial infarction in young males ... 1248
Chest X-ray survey, community-wide.
V. The medical profession [Pamplona] ... 1596
VI. Records and reports [Enterline and Sauer] ... 1613
Childhood, changing causes of death in ... 1246
Children, mental health clinic services for ... 1559
Chloride: *See* Zinc
Cholera:
Burma ... 950, 1016, 1045, 1225, 1262, 1301, 1337, 1377, 1448, 1593
India ... 1016, 1337, 1377, 1485, 1522, 1594, 1684, 1744
India (French) ... 909, 1105, 1225, 1301, 1594
Indochina ... 1378
Pakistan ... 909, 1045, 1105, 1448, 1594, 1715, 1744
World distribution: Asia ... 982, 1133
Civil defense:
And dentists ... 1175
And public health (APHA conference report) ... 1113
Clinic, Central Cooperative, study of follow-up in tuberculosis control. I. Background, objectives, and methodology [Ottenberg, Boucot, Chamberlain, Cohen, and Yelton] ... 853
Clinic services, mental health, for children ... 1559
Committee on Tuberculosis Morbidity Reporting: Report and recommendations, 1951 ... 1293

Page

Conference report:
Southern Branch APHA 1107
State dental directors 1171
Copper: *See* Pentachlorophenates.
Court decisions on municipal milk inspection, recent [Stein and Sonnenshein] 898
Coxiella burnetii in placenta of sheep 1473

D

DDT dusting operations, evaluation of county-wide, in murine typhus control, 1950 [Morlan and Hines] 1052
Death, changing causes of, in childhood [Kahn] 1246
Deaths: Madagascar 881, 908
Dental caries, domestic water and. VII. A study of the fluoride-dental caries relationship in an adult population [Russell and Elvove] 1389
Dental caries experience, effect of topically applied zinc chloride and potassium ferrocyanide on [Anderson and Knutson] 1064
Dental caries reduction and fluorine retention in the bones of white rats 1523
Dental Directors, State, conference report 1171
Dental practice in western Pennsylvania [Ciocco, Altman, and Sissman] 1379
Dentist in atomic warfare 1175
Dependency: *See* Medical
Dermacentor albipictus 1672
Diphtheria: *See* Diseases, communicable: United States.
Disease charts, communicable: United States:
Meningitis, meningococcal 879
Poliomyelitis 879, 1014
Typhoid and paratyphoid fever 1014
Disease incidence: United States 875, 903, 945, 978, 1009, 1039, 1067, 1099, 1127, 1163, 1187, 1219, 1256, 1295, 1332, 1371, 1409, 1443, 1480, 1517, 1553, 1587, 1642, 1677, 1709, 1738
Diseases, communicable: United States: Weekly report by State 877, 906, 948, 980, 1012, 1042, 1071, 1102, 1130, 1165, 1190, 1222, 1259, 1299, 1335, 1375, 1411, 1446, 1483, 1519, 1555, 1591, 1645, 1581, 1712, 1741
Diseases, reported cases:
Canada 880, 908, 982, 1015, 1044, 1073, 1104, 1132, 1167, 1193 1224, 1261, 1300, 1337, 1377, 1413, 1448, 1485, 1521, 1557, 1593, 1647, 1683, 1714, 1743
Cuba 881, 1044, 1224, 1300, 1521, 1714
Finland 982, 1105, 1261, 1413
Jamaica 881, 1167, 1301, 1593
Madagascar 881, 908, 1225, 1743
New Zealand 1105, 1261, 1413, 1683
Norway 1015, 1132, 1193, 1715
Diseases, specific, among males and females 1649
Dogs, histoplasmosis survey of 1533

E

Ectoparasites, rat, and typhus fever in San Antonio, Texas, observations on [Davis] 1717
Elk, winter ticks, and Rocky Mountain spotted fever: A query [Philip and Kohls] 1672

Page

Encephalitis: *See* Diseases, communicable: United States.
Encephalitis in the Missouri River Basin. II. Studies on a focal outbreak, in North Dakota [Wenner, Kamitsuka, Krammer, Cockburn, and Price] 1075
Enterococci detection 1212
Epidemics of influenza and pneumonia, 1918–51 1487
Eskimos, Alaskan, public health problems of 912
Examination for cancer 1339

F

Ferrocyanide, potassium: *See* Zinc chloride and
Filter, membrane, for removing bacteria from liquids 951
Finnish Trambusti tuberculin test 1625
Fluoridation keynoted at State dental conference 1171
Fluoride-dental caries relationship 1389
Fluorides, complex: Caries reduction and fluorine retention in the bones and teeth of white rats [Zipkin and McClure] 1523
Fluorine retention 1523

G

Gland, mammary, Q fever infection in bovine 1685

H

Hagerstown, Md., some conditions leading to medical dependency in 1351
Halogenated derivatives of phenol, toxicity of 1303
Hams, viability of *Brucella melitensis* in naturally infected cured 1402
Hawaii:
  Cage tests with warfarin 1734
  Plague suppressive measures 1727
  Plague surveillance, Hamakua District 1541
Health services, cooperative efforts in (APHA conference report) 1108
Heart disease: Case reports 1248
Helminth parasitism [in Alaska] 928
Histoplasmosis survey of dogs in Louisville, Kentucky [Robinson and Kotcher] 1533
Hospitals, marine, renamed 1138
Hospitals, tuberculosis, nursing in 1272
Hot Springs National Park Cancer Investigation Center streamlined examination 1339
Housing design and home accident prevention, public health considerations on 1461
Housing law enforcement [Johnson] 1451

I

Illness, disabling, from specific causes among males and females of various ages. Sample of white families canvassed at monthly intervals in the Eastern Health District of Baltimore, 1938–43 [Collins, Phillips, and Oliver] 1649
Immunization against tuberculosis, field studies on. I. Tuberculin allergy following BCG vaccination of school children in Muscogee County, Georgia [Shaw] 1415
Indians, Alaskan, public health problems of 912

Page

Industrial sickness absenteeism among males and females during 1950, with index of the previous publications of series [Gafafer] 1550
Infarction, cardiac: *See* Occlusion, coronary.
Infection, *Brucella*, in cured hams 1402
Infection, double, of the rat fleas *X. cheopis* and *N. faciatus* with *Pasteurella* and *Salmonella* [Eskey, Prince, and Fuller] 1318
Influenza: *See* Diseases, communicable: United States.
Influenza, virus, preparation of dried antigen and antiserum for the agglutination-inhibition test for [Hilleman, Buescher, and Smadel] 1195
Influenza and pneumonia, trends and epidemics of, 1918-1951 [Collins and Lehmann] 1487
Insects, Alaska biting, control of [Wilson] 917

K

Keratoconjunctivitis, phlyctenular, among Alaskan Indians and Eskimos [Fritz and Thygeson] 934

L

Law enforcement, housing 1451

M

Mammary gland, Q fever infection in bovine 1685
Man and mice, relative pathogenicity of *Salmonella* strains for 1538
Marine hospitals renamed 1138
Measles: *See* Diseases, communicable: United States.
Mechanisms, body, in tuberculosis 1263
Media, Winter and Sandholzer 1212
Medical care cost, study of. A note on survey methodology [Altman and Wadman] 1019
Medical care in Hagerstown, Md 1351
Medical dependency, some conditions leading to, in Hagerstown, Md. [Lawrence] 1351
Medical social service in a tuberculosis sanatorium [Miller] 987, 1139
Membrane filter in sanitary bacteriology [Clark, Geldreich, Jeter, and Kabler] 951
Meningitis: *See* Diseases, communicable: United States.
Mental health clinic services for children in the United States, 1950 [Pennell, Cameron, and Kramer] 1559
Methods, study, public health 883
Milk inspection, municipal, court decisions 898
Milk sanitation ratings, July 1949–June 1951 1086
Missouri River Basin encephalitis studies 1075
Molluscacides, effect on calves of prolonged oral administration of three potential [Herdt, Loomis, and Nolan] 1313
Morbidity: Baltimore 1227
Morbidity reporting: Tuberculosis 1293
Mortality:
  Childhood 1246
  Neonatal 1038
Mountin, Dr. Joseph W., succeeds Dr. Charles L. Williams as BSS chief 1676
Mus musculus Linn., cage tests with warfarin on *Rattus hawaiiensis* Stone and 1734

N

Page

North Dakota encephalitis outbreak 1075
Nosopsylla fasciatus and *Xenopsylla cheopis*, rat fleas, double infection of, with *Pasteurella* and *Salmonella* 1318
Nurses, public health, 1951 census 1046
Nursing in tuberculosis hospitals [Naylor] 1272

O

Oats, warfarin-treated, use of, as a plague suppressive measure in Hawaii .. 1727
Occlusion, coronary, with myocardial infarction in young males [Levy] 1248
Oral administration of molluscacides 1313

P

Paratyphoid fever: *See* Typhoid and
PAS, effect of, on the emergence of tubercle bacilli resistant to streptomycin 863
Pasteurella and *Salmonella* double infection of the rat fleas with 1318
Pathogen, an unusual enteric [Seligman and Saphra] 1369
Pathogenicity, relative, of certain *Salmonella* strains for man and mice ... 1538
Pennsylvania, western, dental practice 1379
Pentachlorophenates, determination of sodium and copper, in aqueous solutions [Haskins] 1047
Personnel, public health, a national program for training [Tisdale] 1361
Phenol, toxicity of some related halogenated derivatives of 1303
Pittsburgh meeting of Public Health Study Section. Evaluation of study methods 883
Placenta infection, sheep 1473
Plague:
- Belgian Congo 1378
- Brazil 909, 1262, 1594
- Burma 1073, 1378
- Ecuador 1301, 1594
- India 1226, 1485, 1684
- Indochina 882, 909, 1226, 1594
- Madagascar 1226, 1301, 1744
- Peru 1045
- Union of South Africa 1301, 1485, 1744
- Venezuela 1168
- Yemen 1194
- World distribution:
  - Africa 983, 1133
  - Asia 983, 1133
  - South America 983, 1134

Plague infection: United States: Washington 1070, 1482, 1590
Plague in the Territory of Hawaii. II. Plague surveillance, Hamakua District, Island of Hawaii [Gross and Bonnet] 1541
Plague on the high seas [Link] 1466
Plague suppressive measure in Hawaii, use of warfarin-treated oats as a .. 1727
Pneumonia: *See* Diseases, communicable: United States.
Pneumonia trends and epidemics 1487
Poland, BCG vaccinations in 873

Page
Poliomyelitis:
Africa 1168
Europe 1168
United States: *See* Diseases, communicable: United States.
Potassium ferrocyanide: *See* Zinc chloride and
Problems of Alaskan Eskimos, Indians, Aleuts [Haldeman] 912
Public health and mobilization (APHA conference report) 1113
Public health consideration on housing design and home accident prevention [Kent and Pond] 1461
Public health personnel training 1361
Public health problems in Alaska 911, 912, 917, 922, 928, 934, 939, 941
Public health research facilities, Alaska 941
Public Health Service and University of Pittsburgh School of Public Health: Public Health Study Section meeting 883
Public Health Service cooperative investigation on effect of PAS 863
Public Health Service Hospitals: *See* U. S. Marine Hospitals renamed.
Public Health Service publications:
January–June, 1951 1697
July–December, 1950 1091
Public health study methods, evaluation of 883
Publications, industrial sickness absenteeism index of 1552
Publications, Public Health Service 1091, 1697

Q

Q fever in California. IV. Occurrence of *Coxiella burnetii* in the placenta of naturally infected sheep [Welsh, Lennette, Abinanti, and Winn] 1473
Q fever infection in the bovine mammary gland, biochemical study of experimental [Ormsbee] 1685
Q fever outbreak in Sokol, Yugoslavia, August, 1950 [Murray, Djaković, Ljupša, and Snyder 1032

R

Rabies: *See* Diseases, communicable: United States.
Radiography, research in mass. Editorial [Anderson] 851
Rat ectoparasites 1717
Rats, white, fluorine retention in the bones and teeth of 1523
Rattus hawaiiensis Stone and *Mus musculus* Linn., cage tests with warfarin on 1734
Records and reports [in a] community-wide chest X-ray survey 1613
Reports, selected, from Southern Branch, APHA conference 1117
Research, facilities and opportunities for, at the Arctic Health Research Center [Haldeman] 941
Research contributions of BCG vaccination programs. II. Tuberculin sensitivity at different altitudes of residence [Bates, Busk, and Palmer 1427
Rocky Mountain spotted fever: *See* Diseases, communicable: United States; Elk, winter ticks, and

S

Salmonella, double infection of the rat fleas with *Pasteurella* and 1318
Salmonella albany [West and Edwards] 1062
Salmonella california: See Salmonella rostock and
Salmonella homosassa [Edwards and Fife] 1060

Page

Salmonella-like pathogen 1369
Salmonella mendoza: A new *Salmonella* type [Leiguarda, Peso, de Pelazzolo, and Ansiaume] 1478
Salmonella milwaukee [Edwards and Fife] 1059
Salmonella quiniela [Stucker, Galton, Edwards, and Fife] 1058
Salmonella rostock and *Salmonella californa*, changes induced in the flagellar antigens of [Peso and Edwards] 1694
Salmonella strains, relative pathogenicity of, for man and mice [McCullough] 1538
Salmonella thomasville [Edwards, DeCapito, and Fife] 1061
Salmonella types, new 1058, 1478
San Antonio, Texas, rat ectoparasites and typhus fever in 1717
Sanatorium, tuberculosis, medical social service in 987, 1139
Sandholzer media: *See* Winter and
Sanitary engineering graduates, a quantitative study [Lyon] 1177
Sanitary engineers, education and utilization of 1185
Sanitation, water, membrane filter in 951
Scarlet fever: *See* Diseases, communicable: United States.
Serotype, unusual: Type 1296/7 1579
Serotypes, *Shigella boydii* 1327
Sewage and waste disposal projects [Day] 922
Sheep placenta, *Coxiella burnetii* in 1473
Shigella boydii 9, provisional [Ewing, Hucks, and Taylor] 1579
Shigella boydii serotypes, two provisional [Ewing and Taylor] 1327
Sickness, disabling: *See* Illness, disabling.
Sickness absenteeism, industrial 1550
Smallpox:
- Afghanistan 1073
- Algeria 1016, 1744
- Anglo-Egyptian Sudan 1016, 1715
- Bechuanaland 1448
- Belgian Congo 909
- Burma 1194, 1594, 1647, 1744
- Cameroon (British) 1522
- Cameroon (French) 1073, 1226, 1338, 1557
- Ceylon 1338, 1557
- Ecuador 1302, 1486
- Egypt 1448
- French Equatorial Africa 1168, 1262, 1522, 1647, 1684
- French Niger 1194
- French West Africa 950, 1045, 1074, 1168, 1194, 1226, 1262, 1557
- Gold Coast 1226
- India 1045, 1074, 1194, 1378, 1448, 1744
- India (French) 882, 1016, 1045, 1105, 1378, 1486
- Indochina 882, 950, 1106, 1168, 1194, 1262, 1302, 1338, 1486, 1522, 1594, 1684, 1744
- Indonesia 950, 1106, 1168, 1194, 1338, 1449, 1647, 1715
- Iraq 882, 1449
- Ivory Coast 1194
- Korea 909
- Nigeria 1414
- Northern Rhodesia 1486

Page

Smallpox—Continued
Pakistan ... 1074, 1522, 1648
Sierra Leone ... 1106, 1449
Spain ... 1522
Tanganyika ... 950, 1557
Togo (French) ... 1414, 1558, 1594
United States: *See* Diseases, communicable: United States.
Venezuela ... 1715
World distribution:
Africa ... 984, 734
Asia ... 984, 1134
Europe ... 984, 1135
North America ... 1135
South America ... 984, 1135
Social service, medical, in a TB sanatorium ... 987, 1139
Sodium: *See* Pentachlorophenates.
Southern Branch, American Public Health Association, conference report ... 1107
State Dental Directors conference report ... 1171
Streptomycin, effect of PAS on the emergence of tubercle bacilli resistant to ... 863
Study methods, public health ... 883
Survey methodology: Study of medical care cost ... 1019

T

Test, agglutination-inhibition for virus influenza ... 1195
Test, Finnish Trambusti ... 1625
Tests, cage, with warfarin ... 1734
Thrombosis, coronary: *See* Occlusion, coronary.
Ticks, winter, and Rocky Mountain spotted fever ... 1672
Toxicity of some related halogenated derivatives of phenol [Stohlman] ... 1303
Trambusti, Finnish, tuberculin test ... 1625
Trends and epidemics in influenza and pneumonia, 1918–51 ... 1487
Tubercle bacilli resistant to streptomycin, effect of PAS on the emergence of ... 863
Tuberculin allergy following BCG vaccination ... 1415
Tuberculin sensitivity at different altitudes of residence ... 1427
Tuberculin test, Finnish Trambusti. A comparison with the Mantoux test [Edwards and Savonen] ... 1625
Tuberculosis:
Body mechanisms [Payne] ... 1263
Field studies, immunization ... 1415
Laboratory diagnosis courses ... 1170
Medical social service in a sanatorium ... 987, 1139
Morbidity reporting ... 1293
Reportable case ... 1291
Tuberculosis control:
BCG vaccination ... 873, 1158, 1415, 1427
Finnish Trambusti tuberculin test ... 1625
Follow-up study ... 853
X-ray survey, chest ... 1596, 1613
Tuberculosis Control Issue, Public Health Reports, announcement of last ... 1595
Tuberculosis hospitals, nursing in ... 1272
Tuberculosis sanatorium, medical social service in a ... 987, 1139

Page

Tularemia: *See* Diseases, communicable: United States.
Typhoid and paratyphoid fever: *See* Diseases, communicable: United States.
Typhus fever:
Afghanistan ... 1016
Algeria ... 1016, 1262
Ceylon ... 1715
Chile ... 1262, 1594
Ecuador ... 1486
Egypt ... 1226
Eritrea ... 1486, 1594
France ... 1378, 1648
Germany ... 1378, 1414, 1522
India ... 1414, 1594, 1715
Indochina ... 1338, 1414, 1648
Iran ... 1226, 1262, 1338
Iraq ... 1262, 1558, 1684, 1715
Italy ... 1684
Japan ... 1558
Libya ... 1449
Mexico ... 1262, 1716
Morocco (French) ... 1045
New Caledonia ... 1226
Portugal ... 950
Puerto Rico ... 1522
Spain ... 1016, 1414
Tunisia ... 1226
Turkey ... 1262, 1378, 1449, 1558, 1648
United States ... 1052, 1717
Yugoslavia ... 1414
World distribution:
Africa ... 985, 1135
Asia ... 985, 1135
Europe ... 985, 1136
North America ... 985, 1136
South America ... 985, 1136
Typhus fever control, murine ... 1052, 1717

U

U. S. Marine Hospitals renamed ... 1138
University of Pittsburgh Graduate School of Public Health: Public Health Study Section: *See* Public Health Service and

V

Vaccination, BCG ... 873, 1158, 1415, 1427
Virus influenza, test for ... 1195

W

Warfarin, cage tests with, on the Hawaiian rat and the house mouse in Hawaii ... 1734
Warfarin-treated oats as a plague suppressive measure in Hawaii, use of [Gross, Baker, and Bonnet] ... 1727

Page

Water, domestic, and dental caries ---- 1389
Water sanitation, membrane filter in ---- 951
Whooping cough: *See* Diseases, communicable: United States.
Williams, Dr. Charles L. is succeeded by Dr. Joseph W. Mountin as BSS chief ---- 1676
Winter and Sandholzer media and technique, modification of, for enterococci detection [Wang and Dunlop] ---- 1212
World Health Organization BCG vaccination program ---- 873

X

Xenopsylla cheopis and *Nosopsylla fasciatus*, rat fleas, double infection of, with *Pasteurella* and *Salmonella* ---- 1318
X-ray survey, chest ---- 1596, 1613

Y

Yellow fever:
Brazil ---- 950
Cameroon (French) ---- 1106
Colombia ---- 1074, 1194, 1338, 1648
Costa Rica ---- 1106, 1168, 1194, 1262, 1302, 1449, 1594
Ecuador ---- 1486
French West Africa ---- 1302, 1449, 1486, 1558
Gold Coast ---- 882, 909, 950, 1106, 1169, 1194, 1414, 1716, 1558, 1648, 1744
Nigeria ---- 882
Panama ---- 1074
Peru ---- 882, 950
Venezuela ---- 1302, 1522
World distribution:
Africa ---- 986, 1136
North America ---- 986, 1136
South America ---- 986, 1136
Yolk sac technique for isolation of *Brucella* ---- 1204
Yugoslavia Q fever outbreak ---- 1032

Z

Zinc chloride and potassium ferrocyanide, effect of topically applied, on dental caries experience ---- 1064

Author Index

A

Page

Abinanti, Francis R.: *See* Welsh, Hartwell H. (Lennette, Abinanti, and Winn).

Altman, Isidore (and Wadman): Study of the cost of medical care. A note on survey methodology 1019

——— *See also* Ciocco, Antonio (Altman and Sissman).

Anderson, Robert J.: Research in mass radiography. Editorial 851

Anderson, Robert W. (and Knutson): Effect of topically applied zinc chloride and potassium ferrocyanide on dental caries experience 1064

Ansiaume, E. M.: *See* Leiguarda, O. A. (Peso, de Pelazzolo, and Ansiaume).

B

Baker, Robert H.: *See* Gross, Bertram (Baker and Bonnet).

Bates, LeRoy E. (Busk and Palmer): Research contributions of BCG vaccination programs. II. Tuberculin sensitivity at different altitudes of residence 1427

Bonnet, David D. (Mau and Gross): Cage tests with warfarin on the Hawaiian rat, *Rattus hawaiiensis* Stone, and the house mouse, *Mus musculus* Linn., in Hawaii 1734

——— *See also* Gross, Bertram (Baker and Bonnet); Gross, Bertram (and Bonnet).

Boucot, Katherine: *See* Ottenberg, Donald (Boucot, Chamberlain, Cohen, and Yelton).

Buescher, E. L.: *See* Hilleman, M. R. (Buescher and Smadel).

Bunnell, Doris E.: *See* Hutchings, Leslie M. (McCullough, Donham, Eisele, and Bunnell).

Busk, Thøger: *See* Bates, LeRoy E. (Busk and Palmer).

C

Cameron, Dale C.: *See* Pennell, Maryland Y. (Cameron and Kramer).

Chamberlain, W. Edward: *See* Ottenberg, Donald (Boucot, Chamberlain, Cohen, and Yelton).

Ciocco, Antonio (Altman and Sissman): Dental practice in western Pennsylvania 1379

Clark, Harold F. (Geldreich, Jeter, and Kabler): The membrane filter in sanitary bacteriology 951

Cockburn, T. A.: *See* Wenner, H. A. (Kamitsuka, Krammer, Cockburn, and Price).

Cohen, Louis: *See* Ottenberg, Donald (Boucot, Chamberlain, Cohen, and Yelton).

Collins, Selwyn D. (and Lehmann): Trends and epidemics of influenza and pneumonia, 1918–1951 1487

NOTE. Address requests for reprints to the Surgeon General, Public Health Service, Washington 25, D. C.

Page

Collins, Selwyn D. (Phillips and Oliver): Age incidence of specific causes of illness found in monthly canvasses of families—Sample of the Eastern Health District of Baltimore, 1938–43 ---- 1227
——— ——— Disabling illness from specific causes among males and females of various ages. Sample of white families canvassed at monthly intervals in the Eastern Health District of Baltimore, 1938–43 ---- 1649

D

Damon, S. R.: *See* Gay, Kathleen (and Damon).
Davis, David E.: Observations on rat ectoparasites and typhus fever in San Antonio, Texas ---- 1717
Day, E. K.: Sewage and waste disposal problems ---- 922
DeCapito, Thelma: *See* Edwards, P. R. (DeCapito and Fife).
de Pelazzolo, A. Z. R.: *See* Leiguarda, O. A. (Peso, de Pelazzolo, and Ansiaume).
Djaković, Predrag: *See* Murray, Edwards S. (Djaković, Ljupša, and Snyder).
Donham, Charles R.: *See* Hutchings, Leslie M. (McCullough, Donham, Eisele, and Bunnell).
Dunlop, Stuart G.: *See* Wang, Wen-Lan Lou (and Dunlop).

E

Edwards, P. R. (DeCapito and Fife): *Salmonella thomasville* ---- 1061
——— (and Fife): *Salmonella homosassa* ---- 1060
——— ——— *Salmonella milwaukee* ---- 1059
——— *See also* Peso, O. A. (and Edwards); Stucker, Calvin L. (Galton, Edwards, and Fife); West, Mary G. (and Edwards).
Edwards, Phyllis Q. (and Savonen): The Finnish Trambusti tuberculin test—A comparison with the Mantoux test ---- 1625
Eisele, C. Wesley: *See* Hutchings, Leslie M. (McCullough, Donham, Eisele, and Bunnell).
Elvove, Elias: *See* Russell, A. L. (and Elvove).
Enterline, Philip E. (and Sauer): Community-wide chest X-ray survey. VI. Records and reports ---- 1613
Eskey, C. R. (Prince and Fuller): Double infection of the rat fleas, *X. cheopis* and *N. fasciatus* with *Pasteurella* and *Salmonella* ---- 1318
Ewing, W. H. (Hucks and Taylor): Provisional *Shigella boydii* 9 ---- 1579
——— (and Taylor): Two provisional *Shigella boydii* serotypes ---- 1327

F

Fife, Mary A.: *See* Edwards, P. R. (DeCapito and Fife); Edwards, P. R. (and Fife); Stucker, Calvin L. (Galton, Edwards, and Fife).
Fritz, Milo H. (and Thygeson): Phlyctenular keratoconjunctivitis among Alaskan Indians and Eskimos ---- 937
Fuller, Frank B.: *See* Eskey, C. R. (Prince and Fuller).

G

Gafafer, W. M.: Industrial sickness absenteeism among males and females during 1950—With index of the previous publications of the series ---- 1550
Galton, Mildred M.: *See* Stucker, Calvin L. (Galton, Edwards, and Fife).
Gay, Kathleen (and Damon): A yolk sac technique for the routine isolation of *Brucella*. Injection of clotted blood specimens into embryonating eggs with recovery of all three species ---- 1204

Page

Geldreich, Edwin E.: *See* Clark, Harold F. (Geldreich, Jeter, and Kabler).
Gross, Bertram (Baker and Bonnet): Use of warfarin-treated oats as a plague suppressive measure in Hawaii 1727
———(and Bonnet): Plague in the Territory of Hawaii. II. Plague surveillance, Hamakua District, Island of Hawaii 1541
———*See also* Bonnet, David D. (Mau and Gross).

H

Haldeman, Jack C.: Facilities and opportunities for research at the Arctic Health Research Center 941
———Problems of Alaskan Eskimos, Indians, Aleuts 912
Haskins, W. T.: Determination of sodium and copper pentachlorophenates in dilute aqueous solutions 1047
Herdt, Jean R. (Loomis and Nolan): Effect on calves of prolonged oral administration of three potential molluscacides 1313
Hilleman, M. R. (Buescher and Smadel): Preparation of dried antigen and antiserum for the agglutination-inhibition test for virus influenza 1195
Hines, Virginia D.: *See* Morlan, Harvey B. (and Hines).
Hucks, M. C.: *See* Ewing, W. H. (Hucks and Taylor).
Hutchings, Leslie M. (McCullough, Donham, Eisele, and Bunnell): The viability of *Brucella melitensis* in naturally infected cured hams 1402

I

Irving, Laurence: Climatic adaptation in arctic and tropic animals 939

J

Jeter, Harold L.: *See* Clark, Harold F. (Geldreich, Jeter, and Kabler).
Johnson, Ralph J.: Housing law enforcement 1451

K

Kabler, Paul W.: *See* Clark, Harold F. (Geldreich, Jeter, and Kabler).
Kahn, Harold A.: Changing causes of death in childhood 1246
Kamitsuka, Paul.: *See* Wenner, H. A. (Kamitsuka, Krammer, Cockburn, and Price).
Kent, Frederick S. (and Pond): Public health consideration on housing design and home accident prevention 1461
Knutson, John W.: *See* Anderson, Robert W. (and Knutson).
Kohls, Glen M.: *See* Philip, Cornelius B. (and Kohls).
Koplin, Allen N.: Experience with a streamlined examination—Hot Springs National Park Cancer Investigation Center 1339
Kotcher, Emil: *See* Robinson, John W. (and Kotcher).
Kramer, Morton: *See* Pennell, Maryland Y. (Cameron and Kramer).
Krammer, M. C.: *See* Wenner, H. A. (Kamitsuka, Krammer, Cockburn, and Price).

L

Lawrence, P. S.: Some conditions leading to medical dependency in Hagerstown, Md 1351
Lehmann, Josephine: *See* Collins, Selwyn D. (and Lehmann).
Leiguarda, R. H. (Peso, de Pelazzolo, and Ansiaume): *Salmonella mendoza:* A new *Salmonella* type 1478
Lennette, Edwin H.: *See* Welsh, Hartwell H. (Lennette, Abinanti, and Winn).

Page

Levy, Tracy: Coronary occlusion with myocardial infarction in young males. Report of four cases 1248
Link, Vernon B.: Plague on the high seas 1466
Ljupša, Franja: *See* Murray, Edward S. (Djaković, Ljupša, and Snyder).
Loomis, Ladd N.: *See* Herdt, Jean R. (Loomis and Nolan).
Lyon, Walter A.: A quantitative study of sanitary engineering graduates. 1177

M

Mau, Edward S. C.: *See* Bonnet, Edward D. (Mau and Gross).
McClure, F. J.: *See* Zipkin, I. (and McClure).
McCullough, Norman B.: Relative pathogenicity of certain *Salmonella* strains for man and mice 1538
———*See also* Hutchings, Leslie M. (McCullough, Donham, Eisele, and Bunnell).
Miller, Pauline: Medical social service in a tuberculosis sanatorium [Parts I and II] 987, 1139
Morlan, Harvey B. (and Hines): Evaluation of county-wide DDT dusting operations in murine typhus control, 1950 1052
Murray, Edward S. (Djaković, Ljupša, and Snyder): An outbreak of Q fever in Sokol, Yugoslavia, August 1950 1032

N

Naylor, Martha Ball: Nursing in tuberculosis hospitals 1272
Nolan, M. O.: *See* Herdt, Jean R. (Loomis and Nolan).

O

Oliver, Dorothy S.: *See* Collins, Selwyn D. (Phillips and Oliver).
Ormsbee, Richard A.: A biochemical study of experimental Q fever infection in the bovine mammary gland 1685
Ottenberg, Donald (Boucot, Chamberlain, Cohen, and Yelton): Central cooperative clinic study of follow-up in tuberculosis control. I. Background, objectives, and methodology 853

P

Palmer, Carroll E.: *See* Bates, LeRoy E. (Busk and Palmer).
Pamplona, Paul A.: Community-wide chest X-ray survey. V. The medical profession 1596
Payne, Howard M.: Body mechanisms in progressive tuberculosis 1263
Pennell, Maryland Y. (Cameron and Kramer): Mental health clinic services for children in the United States, 1950 1559
Peso, O. A. (and Edwards): Changes induced in the flagellar antigens of *Salmonella rostock* and *Salmonella california* 1694
——— *See also* Leiguarda, O. A. (Peso, de Pelazzolo, and Ansiaume).
Philip, Cornelius B. (and Kohls): Elk, winter ticks, and Rocky Mountain spotted fever: A query 1672
Phillips, F. Ruth: *See* Collins, Selwyn D. (Phillips and Oliver).
Pond, M. Allen: *See* Kent, Frederick S. (and Pond).
Price, E. R.: *See* Wenner, H. A. (Kamitsuka, Krammer, Cockburn, and Price).
Prince, Frank M.: *See* Eskey, C. R. (Prince and Fuller).

R

Rausch, Robert: Biotic interrelationships of helminth parasitism 928

Page

Robinson, John W. (and Kotcher): Histoplasmosis survey of dogs in Louisville, Kentucky 1533
Russell, A. L. (and Elvove): Domestic water and dental caries. VII. A study of the fluoride-dental caries relationship in an adult population .. 1389

S

Saphra, Ivan: *See* Seligman, Erich (and Saphra).
Sauer, Herbert I.: *See* Enterline, Philip E. (and Sauer).
Savonen, Severi: *See* Edwards, Phyllis Q. (and Savonen).
Seligman, Erich (and Saphra): An unusual enteric pathogen 1369
Shaw, Lawrence W.: Field studies on immunization against tuberculosis. I. Tuberculin allergy following BCG vaccination of school children in Muscogee County, Georgia 1415
Sissman, Isaac: *See* Ciocco, Antonio (Altman and Sissman).
Smadel, J. E.: *See* Hilleman, M. R. (Buescher and Smadel).
Snyder, John C.: *See* Murray, Edward S. (Djaković, Ljupša, and Snyder).
Sonenshein, Israel L.: *See* Stein, Murray (and Sonenshein).
Stein, Murray (and Sonenshein): Recent court decisions on municipal milk inspection 898
Stohlman, E. F.: The toxicity of some related halogenated derivatives of phenol 1303
Stucker, Calvin L. (Galton, Edwards, and Fife): *Salmonella quiniela* 1058

T

Taylor, M. W.: *See* Ewing, W. H. (Hucks and Taylor); Ewing, W. H. (and Taylor).
Thomas, Evan W.: The influence of modern treatment on syphilis control .. 1573
Thygeson, Phillips: *See* Fritz, Milo H. (and Thygeson).
Tisdale, Ellis S.: A national program for training public health personnel .. 1361

W

Wadman, Ruth: *See* Altman, Isidore (and Wadman).
Wang, Wen-Lan Lou (and Dunlop): A modification of the Winter and Sandholzer media and technique for enterococci detection 1212
Welsh, Hartwell H. (Lennette, Abinanti, and Winn): Q fever in California. IV. Occurrence of *Coxiella burnetii* in the placenta of naturally infected sheep 1473
Wenner, H. A. (Kamitsuka, Krammer, Cockburn, and Price): Encephalitis in the Missouri River Basin. II. Studies on a focal outbreak of encephalitis in North Dakota 1075
West, Mary G. (and Edwards): *Salomella albany* 1062
Wilson, Charles S.: Control of Alaskan biting insects 917
Winn, John F.: *See* Welsh, Hartwell H. (Lennette, Abinanti, and Winn).

Y

Yelton, Sara E.: *See* Ottenberg, Donald (Boucot, Chamberlain, Cohen, and Yelton).

Z

Zipkin, I. (and McClure): Complex fluorides: Caries reduction and fluorine retention in the bones and teeth of white rats 1523

U. S. GOVERNMENT PRINTING OFFICE: 1952